arXiv:1607.02719v1 [math.RA] 10 Jul 2016

# Quadratic descent of totally decomposable orthogonal involutions in characteristic two

A.-H. Nokhodkar

July 12, 2016

**Abstract**

We investigate the quadratic descent of totally decomposable algebras with involution of orthogonal type in characteristic two. Both separable and inseparable extensions are included.
*Mathematics Subject Classification:* 16W10, 16K20, 11E39.

## 1 Introduction

Let $K/F$ be a field extension and let $(A,\sigma)$ be a central simple algebra with involution over $K$. By a *descent* of $(A,\sigma)$ to $F$ we mean an $F$-algebra with involution $(B,\sigma')$ satisfying $(A,\sigma)\simeq_K (B,\sigma')\otimes_F (K,\mathrm{id})$. In the case where $\mathrm{char}\, F\neq 2$ and $K/F$ is a quadratic extension, the descent of orthogonal involutions of degree 2 and 4 over $K$ was studied in [2] and necessary and sufficient conditions in terms of the discriminant and the Clifford algebra were given. Also, if $\mathrm{char}\, F=2$ and $K/F$ is a finite extension satisfying $K^2\subseteq F$, a criterion for totally decomposable algebras with orthogonal involution over $K$ to have a descent to $F$ was obtained in [7, (6.2)].

Let $(A,\sigma)$ be a totally decomposable algebra with orthogonal involution over a field $F$ of characteristic 2. In [7], a maximal commutative Frobenius subalgebra $\Phi(A,\sigma)$ of $A$ was introduced and the relation between this subalgebra and the *Pfister invariant* $\mathfrak{Pf}(A,\sigma)$, a bilinear Pfister form associated to $(A,\sigma)$ defined in [3], was studied. It was also shown that totally decomposable orthogonal involutions can be classified, up to conjugation, by their Pfister invariant (see [7, (6.5)]).

In this work we study the quadratic descent of totally decomposable orthogonal involutions in characteristic 2. We first consider inseparable extensions. If $K=F(\sqrt{\alpha})$ is a quadratic extension of $F$ and $(A,\sigma)$ is a totally decomposable algebra with orthogonal involution over $K$, we find necessary and sufficient conditions for $(A,\sigma)$, in terms of $\Phi(A,\sigma)$ and $\mathfrak{Pf}(A,\sigma)$, to has a descent to $F$. It is also shown that if $(A,\sigma)$ has a descent to $F$, then it has a *totally decomposable descent* to $F$, i.e., a descent which is totally decomposable itself (see (3.4)). This result complements [7, (6.2)] for quadratic extensions.

We next consider separable extensions. In §4 we investigate the quadratic descent of quaternion algebras with orthogonal involution. In §5 some properties of totally singular conic algebras are studied. These results will be used in §6 to obtain a descent property of the Pfister invariant. Finally, in §7 we state our

main result (7.3) which asserts that for a separable quadratic extension $K/F$ and a totally decomposable algebra with orthogonal involution $(A,\sigma)$ over $K$, the following conditions are equivalent: (1) $(A,\sigma)$ has a descent to $F$. (2) $(A,\sigma)$ has a totally decomposable descent to $F$. (3) The corestriction $\mathrm{cor}_{K/F}(A)$ splits and $\mathfrak{Pf}(A,\sigma) \simeq \mathfrak{b}_K$ for some symmetric bilinear form $\mathfrak{b}$ over $F$. In the case where $\sigma$ is isotropic, these conditions are also equivalent to: (4) $\mathrm{cor}_{K/F}(A)$ splits and $\Phi(A,\sigma) \simeq S \otimes_F K$ for some totally singular conic $F$-algebra $S$.

# 2 Preliminaries

Throughout this work, all fields are supposed to be of characteristic 2.

Let $A$ be a central simple algebra over a field $F$. An *involution* on $A$ is a map $\sigma : A \to A$ satisfying $\sigma^2(x) = x$, $\sigma(x+y) = \sigma(x) + \sigma(y)$ and $\sigma(xy) = \sigma(y)\sigma(x)$ for $x, y \in A$. For an algebra with involution $(A,\sigma)$ we define the subspaces

$$\mathrm{Alt}(A,\sigma) = \{x - \sigma(x) \mid x \in A\} \quad \text{and} \quad \mathrm{Sym}(A,\sigma) = \{x \in A \mid \sigma(x) = x\}.$$

If $\sigma|_F = \mathrm{id}$, we say that $\sigma$ is of *the first kind*. Otherwise, $\sigma$ is said to be of *the second kind*. An involution $\sigma$ of the first kind is called *symplectic* if after scalar extension to a splitting field, it becomes adjoint to an alternating bilinear form. Otherwise, it is called *orthogonal*. According to [5, (2.6 (2)], $\sigma$ is orthogonal if and only if $1 \notin \mathrm{Alt}(A,\sigma)$. We denote the discriminant of an orthogonal involution $\sigma$ by $\mathrm{disc}\,\sigma$ (see [5, (7.2)]). An algebra with involution $(A,\sigma)$ (or the involution $\sigma$ itself) is called *isotropic* if there exists a nonzero element $x \in A$ such that $\sigma(x)x = 0$. Otherwise, it is called *anisotropic*.

A *quaternion algebra* over a field $F$ is a four-dimensional central simple $F$-algebra. Every quaternion algebra $Q$ has a basis $(1,u,v,w)$, called a *quaternion basis*, satisfying $u^2 + u \in F$, $v^2 \in F^\times$ and $w = uv = vu + v$ (see [5, p. 25]). In this case, if $a = u^2 + u \in F$ and $b = v^2 \in F^\times$, then $Q$ is denoted by $[a,b)_F$. If $b \in F^{\times 2}$ or $a \in \wp(F) := \{x^2 + x \mid x \in F\}$, then $Q$ splits. Also, it is readily verified that every element $v' \in Q \setminus F$ satisfying $v'^2 \in F^\times$ extends to a quaternion basis $(1,u',v',w')$ of $Q$.

Let $V$ be a finite-dimensional vector space over a field $F$. A symmetric bilinear form $\mathfrak{b} : V \times V \to F$ is called *metabolic* if there exists a subspace $W$ of $V$ with $\dim_F W = \frac{1}{2}\dim_F V$ such that $\mathfrak{b}|_{W\times W} = 0$. We say that two bilinear forms $\mathfrak{b}$ and $\mathfrak{b}'$ are *similar* if $\mathfrak{b} \simeq \lambda\mathfrak{b}'$ for some $\lambda \in F^\times$. For $\alpha \in F$ the isometry class of the symmetric bilinear form $\mathfrak{b}((x_1,x_2),(y_1,y_2)) = x_1y_1 + \alpha x_2y_2$ on $F^2$ is denoted by $\langle 1,\alpha\rangle$. Also, for $\alpha_1,\cdots,\alpha_n \in F^\times$, the form $\langle 1,\alpha_1\rangle \otimes \cdots \otimes \langle 1,\alpha_n\rangle$ is called a *bilinear (n-fold) Pfister form* and is denoted by $\langle\langle \alpha_1,\cdots,\alpha_n\rangle\rangle$. Finally, if $K/F$ is a field extension and $\mathfrak{b}$ is a bilinear form over $F$, the scalar extension of $\mathfrak{b}$ to $K$ is denoted by $\mathfrak{b}_K$.

# 3 Inseparable descent

We begin our discussion with a definition from [7].

**Definition 3.1.** An algebra $A$ over a field $F$ is called a *totally singular conic algebra* if $x^2 \in F$ for every $x \in R$.

According to [7, (3.2)], every totally singular conic algebra is a local commutative algebra.

**Definition 3.2.** An algebra with involution $(A, \sigma)$ over a field $F$ is called *totally decomposable* if it is isomorphic to a tensor product of quaternion $F$-algebras with involution.

Note that if $\sigma$ is orthogonal and $(A, \sigma) \simeq \bigotimes_{i=1}^{n} (Q_i, \sigma_i)$ is a decomposition of $(A, \sigma)$ into quaternion $F$-algebras with involution, then every $\sigma_i$ is necessarily orthogonal by [5, (2.21)].

Let $(A, \sigma)$ be a totally decomposable algebra of degree $2^n$ with orthogonal involution over $F$. In [7], it was shown that there exists a $2^n$-dimensional totally singular conic $F$-algebra $\Phi(A, \sigma) \subseteq F \oplus \mathrm{Alt}(A, \sigma)$ which is maximal commutative (i.e., $C_A(\Phi(A, \sigma)) = \Phi(A, \sigma)$) and is generated, as an $F$-algebra, by $n$ elements. Also, according to [7, (5.10)] the subalgebra $\Phi(A, \sigma)$ is uniquely determined, up to isomorphism. By [7, (4.6)] we have $\Phi(A, \sigma) = F[v_1, \cdots, v_n]$ for some $v_1, \cdots, v_n \in \mathrm{Alt}(A, \sigma)$.

**Lemma 3.3.** *Let $K/F$ be a finite field extension satisfying $K^2 \subseteq F$ and let $(Q, \sigma)$ be a quaternion algebra with orthogonal involution over $K$. For every $v \in \mathrm{Alt}(Q, \sigma)$ satisfying $v^2 \in F^\times$, there exists a descent $(Q_0, \sigma_0)$ of $(Q, \sigma)$ to $F$ such that $v \in \mathrm{Alt}(Q_0, \sigma_0)$.*

*Proof.* See [7, (6.1)]. □

**Lemma 3.4.** *Let $K/F$ be a finite field extension satisfying $K^2 \subseteq F$ and let $(A, \sigma)$ be a totally decomposable algebra with orthogonal involution over $K$. Then $(A, \sigma)$ has a totally decomposable descent to $F$ if and only if $x^2 \in F$ for every $x \in \Phi(A, \sigma)$.*

*Proof.* If $x^2 \in F$ for every $x \in \Phi(A, \sigma)$, then the proof of [7, (6.2)] shows that $(A, \sigma)$ has a totally decomposable descent to $F$. Conversely, if $(B, \sigma')$ is a totally decomposable descent of $(A, \sigma)$ to $F$, then $\Phi(A, \sigma) \simeq \Phi(B, \sigma') \otimes_F K$ as $K$-algebras. Since $K^2 \subseteq F$ and $v^2 \in F$ for every $v \in \Phi(B, \sigma')$, we have $x^2 \in F$ for every $x \in \Phi(A, \sigma)$. □

**Definition 3.5.** Let $(A, \sigma) \simeq \bigotimes_{i=1}^{n} (Q_i, \sigma_i)$ be a totally decomposable algebra with orthogonal involution over a field $F$ and let $\alpha_i \in F^\times$ be a representative of the class $\operatorname{disc} \sigma_i \in F^\times / F^{\times 2}$. As in [3], we call the bilinear Pfister form $\langle\langle \alpha_1, \cdots, \alpha_n \rangle\rangle$, the *Pfister invariant* of $(A, \sigma)$ and we denote it by $\mathfrak{Pf}(A, \sigma)$.

Note that according to [3, (7.5)], the isometry class of $\mathfrak{Pf}(A, \sigma)$ does not depend on the decomposition of $(A, \sigma)$. Also, as observed in [7, (5.5)], the algebra $\Phi(A, \sigma)$ can be considered as an underlying vector space of $\mathfrak{Pf}(A, \sigma)$ in such a way that for $v, w \in \Phi(A, \sigma)$, $\mathfrak{Pf}(A, \sigma)(v, w)$ is the unique element $\alpha \in F$ satisfying $vw + \alpha \in \mathrm{Alt}(A, \sigma)$. In particular, $\mathfrak{Pf}(A, \sigma)(v, v) = v^2$ for every $v \in \Phi(A, \sigma)$.

The next result complements [7, (6.2)] for quadratic extensions.

**Theorem 3.6.** *Let $K = F(\sqrt{\alpha})$ be a quadratic extension of a field $F$. For a totally decomposable algebra with orthogonal involution $(A, \sigma)$ over $K$ the following conditions are equivalent:*

*(1) $(A, \sigma)$ has a descent to $F$.*

*(2) $(A, \sigma)$ has a totally decomposable descent to $F$.*

*(3)* $x^2 \in F$ *for every* $x \in \Phi(A,\sigma)$.

*(4)* $\mathfrak{Pf}(A,\sigma) \simeq \mathfrak{b}_K$ *for some symmetric bilinear form* $\mathfrak{b}$ *over* $F$.

*Proof.* The equivalence of (2) and (3) follows from (3.4). The implication (2) $\Rightarrow$ (1) is evident and (2) $\Rightarrow$ (4) follows by setting $\mathfrak{b} = \mathfrak{Pf}(C,\tau)$, where $(C,\tau)$ is a totally decomposable descent of $(A,\sigma)$. It suffices therefore to prove the implications (1) $\Rightarrow$ (3) and (4) $\Rightarrow$ (3).

(1) $\Rightarrow$ (3): Let $(B,\sigma')$ be a descent of $(A,\sigma)$ and let

$$g : (A,\sigma) \xrightarrow{\sim} (B,\sigma') \otimes_F (K, \mathrm{id}),$$

be an isomorphism of $K$-algebras with involution. Write $\Phi(A,\sigma) = F[v_1,\cdots,v_n]$ for some $v_1,\cdots,v_n \in \mathrm{Alt}(A,\sigma)$. Since $g(\mathrm{Alt}(A,\sigma)) = \mathrm{Alt}(B,\sigma') \otimes_F K$, there exist $u_i, w_i \in \mathrm{Alt}(B,\sigma')$ such that $g(v_i) = u_i \otimes 1 + w_i \otimes \sqrt{\alpha}$, $i = 1,\cdots,n$. Hence

$$g(v_i)^2 = (u_i^2 + \alpha w_i^2) \otimes 1 + (u_i w_i + w_i u_i) \otimes \sqrt{\alpha}.$$

Since $v_i^2 \in K$, the element $g(v_i)^2$ lies in the center of $B \otimes_F K$, i.e., $u_i^2 + \alpha w_i^2 \in F$ and $u_i w_i + w_i u_i \in F$. We have $u_i w_i + w_i u_i = u_i w_i - \sigma(u_i w_i) \in \mathrm{Alt}(B,\sigma')$. The orthogonality of $\sigma'$ implies that $1 \notin \mathrm{Alt}(B,\sigma')$, hence $u_i w_i + w_i u_i = 0$. It follows that $g(v_i)^2 = (u_i^2 + \alpha w_i^2) \otimes 1 \in F \otimes_F F$, i.e., $v_i^2 \in F$. Since $K^2 \subseteq F$ and $\Phi(A,\sigma)$ is commutative, we get $x^2 \in F$ for every $x \in \Phi(A,\sigma)$.

(4) $\Rightarrow$ (3): Let $V$ be an underlying vector space of $\mathfrak{b}$ and consider an isometry

$$h : (\Phi(A,\sigma), \mathfrak{Pf}(A,\sigma)) \xrightarrow{\sim} (V \otimes_F K, \mathfrak{b}_K).$$

Let $v \in \Phi(A,\sigma)$ and write $h(v) = u \otimes 1 + w \otimes \sqrt{\alpha}$ for some $u, w \in V$. Then

$$\begin{aligned} v^2 &= \mathfrak{Pf}(A,\sigma)(v,v) = \mathfrak{b}_K(h(v),h(v)) = \mathfrak{b}_K(u \otimes 1 + w \otimes \sqrt{\alpha}, u \otimes 1 + w \otimes \sqrt{\alpha}) \\ &= \mathfrak{b}(u,u) + \sqrt{\alpha}\mathfrak{b}(u,w) + \sqrt{\alpha}\mathfrak{b}(w,u) + \alpha\mathfrak{b}(w,w) = \mathfrak{b}(u,u) + \alpha\mathfrak{b}(w,w) \in F, \end{aligned}$$

which completes the proof. □

# 4 Separable descent of quaternion algebras

From now on we fix a separable quadratic extension $K/F$ and two elements $\eta \in K$ and $\delta \in F \setminus \wp(F)$ such that $K = F(\eta)$ and $\eta^2 + \eta = \delta$. The *trace* of an element $x \in K$ over $F$ is denoted by $T_{K/F}(x)$. Hence $T_{K/F}(a + b\eta) = b$ for $a, b \in F$.

Let $A$ be a central simple algebra over $K$ and let $\iota$ be the nontrivial automorphism of $K/F$. Define the *conjugate* algebra ${}^\iota A = \{{}^\iota x \mid x \in A\}$ with the operations

$${}^\iota x + {}^\iota y = {}^\iota(x+y), \quad {}^\iota x {}^\iota y = {}^\iota(xy) \quad \text{and} \quad {}^\iota(\alpha x) = \iota(\alpha) {}^\iota x,$$

for $x, y \in A$ and $\alpha \in K$. Let $s : {}^\iota A \otimes_K A \to {}^\iota A \otimes_K A$ be the switch map induced by $s({}^\iota x \otimes y) = {}^\iota y \otimes x$ for $x, y \in A$. The *corestriction* of $A$ is defined as follows:

$$\mathrm{cor}_{K/F}(A) = \{u \in {}^\iota A \otimes_K A \mid s(u) = u\}.$$

According to [5, (3.13 (4))], $\mathrm{cor}_{K/F}(A)$ is a central simple algebra over $F$.

**Lemma 4.1.** *A quaternion $K$-algebra $Q$ has a descent to $F$ if and only if* $\operatorname{cor}_{K/F}(Q)$ *splits.*

*Proof.* The result follows from [5, (2.22)] and [9, Ch. 8, (9.5)]. □

The following result can be found in [5, Ch. I, Exercise 9].

**Lemma 4.2.** *If $a \in K$ and $b \in F^\times$, then* $\operatorname{cor}_{K/F}([a,b)_K) \sim [T_{K/F}(a), b)_F$, *where $\sim$ denotes Brauer-equivalence.*

**Lemma 4.3.** *Let $Q = [a,b)_E$ be a quaternion algebra over a field $E$. If $b \notin E^{\times 2}$, then $Q$ splits if and only if $a + c^2 b \in \wp(E)$ for some $c \in E$.*

*Proof.* Consider the quadratic étale extension $E_a := E[X]/(X^2 + X + a)$ of $E$. By [4, (98.14) (5)], $Q$ splits if and only if $b$ is a norm in $E_a$, or equivalently $b = \beta^2 + \beta\lambda + \lambda^2 a$ for some $\beta, \lambda \in E$. As $b \notin E^{\times 2}$, we have $\lambda \neq 0$. Hence $Q$ splits if and only if $a + \lambda^{-2} b = \lambda^{-1}\beta + (\lambda^{-1}\beta)^2 \in \wp(E)$. □

**Corollary 4.4.** *Let $[a,b)_E$ and $[c,b)_E$ be quaternion algebras over a field $E$. If $b \notin E^{\times 2}$, then $[a,b)_E \simeq [c,b)_E$ if and only if $a + c + d^2 b \in \wp(E)$ for some $d \in E$.*

*Proof.* The result follows from (4.3) and [9, Ch. 8, (11.1)]. □

**Lemma 4.5.** *Let $(Q,\sigma)$ be a quaternion algebra with orthogonal involution over a field $E$ and let $(1,u,v,w)$ be a quaternion basis of $Q$. If $v \in \operatorname{Alt}(Q,\sigma)$, then $\sigma(u) = u$.*

*Proof.* Since $\sigma(u) - u \in \operatorname{Alt}(Q,\sigma)$ and $\dim_E \operatorname{Alt}(Q,\sigma) = 1$, there exists $a \in E$ such that $\sigma(u) - u = av$. Hence $\sigma(uv) = v(u + av) = uv + v + ab$, where $b = v^2 \in F^\times$. It follows that $v + ab \in \operatorname{Alt}(Q,\sigma)$, i.e., $ab \in \operatorname{Alt}(Q,\sigma)$. As $\sigma$ is orthogonal, we get $a = 0$. □

The next result is easily deduced from [5, (7.4)].

**Lemma 4.6.** *Let $E$ be a field and let $t$ be the transpose involution on $M_2(E)$. For a quaternion algebra with orthogonal involution $(Q,\sigma)$ over $E$ we have $(Q,\sigma) \simeq (M_2(E), t)$ if and only if* $\operatorname{disc}\sigma$ *is trivial.*

**Proposition 4.7.** *(Compare [2, (2.4)]) A quaternion algebra with orthogonal involution $(Q,\sigma)$ over $K$ has a descent to $F$ if and only if* $\operatorname{cor}_{K/F}(Q)$ *splits and there exists $v \in \operatorname{Alt}(Q,\sigma)$ such that $v^2 \in F^\times$. In addition, if $(Q_0,\sigma_0)$ is a descent of $(Q,\sigma)$ to $F$, then $v \in \operatorname{Alt}(Q_0,\sigma_0)$.*

*Proof.* If $Q \simeq_K Q_0 \otimes_F K$ for some $\sigma$-invariant quaternion $F$-algebra $Q_0 \subseteq Q$, then $\operatorname{cor}_{K/F}(Q)$ splits by (4.1). Also for every unit $v \in \operatorname{Alt}(Q_0, \sigma|_{Q_0})$ we have $v^2 \in F^\times$. Conversely, let $b = v^2 \in F^\times$. If $b \in F^{\times 2}$, then by (4.6) we have

$$(Q,\sigma) \simeq_K (M_2(K), t) \simeq_K (M_2(F), t) \otimes_F (K, \mathrm{id}),$$

and we are done. Suppose that $b \notin F^{\times 2}$. Extend $v$ to a quaternion basis $(1,u,v,w)$ of $Q$ and set $a = u^2 + u \in K$. Then $Q \simeq [a,b)_K$. Write $a = \beta + \lambda\eta$ for some $\beta, \lambda \in F$. By (4.2), $\operatorname{cor}_{K/F}(Q) \sim [T_{K/F}(a), b)_F$, thus $[\lambda, b)_F$ splits. By (4.3) there exist $d, e \in F$ such that $\lambda = d^2 + d + e^2 b$. Set $u' = u + e\eta v + d\eta \in Q$ and $c = \beta + d^2\delta + e^2\delta b \in F$. Then $u'v + vu' = v$ and $u'^2 + u' = c$. Hence

the $F$-algebra $Q_0$ generated by $u'$ and $v$ is a quaternion algebra, isomorphic to $[c,b)_F$. We also have

$$\begin{aligned} a+c &= (\beta+\lambda\eta)+(\beta+d^2\delta+e^2\delta b) = (d^2+d+e^2b)\eta+d^2\delta+e^2\delta b \\ &= d^2\eta+d^2\delta+d\eta+e^2\eta b+e^2\delta b = (d\eta)^2+d\eta+(e\eta)^2 b. \end{aligned}$$

Hence $a+c+(e\eta)^2 b \in \wp(K)$. Using (4.4) we get $[c,b)_K \simeq [a,b)_K$, i.e., $Q \simeq_K Q_0 \otimes_F K$. Note that $(1,u',v,u'v)$ is a quaternion basis of $[c,b)_K$, hence $\sigma(u') = u'$ by (4.5). It follows that $\sigma(Q_0) = Q_0$, i.e., $(Q,\sigma) \simeq_K (Q_0, \sigma|_{Q_0}) \otimes_F (K, \mathrm{id})$.

To prove the last assertion of the result, observe that as $\mathrm{Alt}(Q,\sigma)$ is one-dimensional, for every unit $w \in \mathrm{Alt}(Q_0,\sigma_0)$ we have $\mathrm{Alt}(Q,\sigma) = Kw$. Hence one can write $v = \alpha w$ for some $\alpha \in K^\times$. Since $v^2, w^2 \in F^\times$ and $K^{\times 2} \cap F^\times = F^{\times 2}$, we get $\alpha \in F^\times$, hence $v \in \mathrm{Alt}(Q_0,\sigma_0)$. □

# 5 Some descent properties of totally singular conic algebras

We recall that $K = F(\eta)$ is a separable quadratic extension of $F$ with $\eta^2+\eta = \delta \in F \setminus \wp(F)$.

**Lemma 5.1.** *Let $(A,\sigma)$ be a totally decomposable algebra with orthogonal involution over $K$. Suppose that $\mathfrak{Pf}(A,\sigma) \simeq \langle\langle \alpha_1, \cdots, \alpha_n \rangle\rangle$ for some $\alpha_1, \cdots, \alpha_n \in K^\times$. If $(A,\sigma)$ has a descent to $F$, then $\mathfrak{Pf}(A,\sigma) \simeq \langle\langle \iota(\alpha_1), \cdots, \iota(\alpha_n) \rangle\rangle$, where $\iota$ is the nontrivial automorphism of $K/F$.*

*Proof.* Let $(B,\sigma')$ be a descent of $(A,\sigma)$ to $F$ and consider an isomorphism of $K$-algebras with involution

$$f : (A,\sigma) \xrightarrow{\sim} (B,\sigma') \otimes_F (K, \mathrm{id}).$$

For $x \in B$ and $\alpha \in K$, the assignment $x \otimes \alpha \mapsto x \otimes \iota(\alpha)$ induces an $F$-algebra isomorphism $g : B \otimes_F K \to B \otimes_F K$ satisfying $g \circ (\sigma' \otimes \mathrm{id}) = (\sigma' \otimes \mathrm{id}) \circ g$. The composition

$$h = f^{-1} \circ g \circ f : A \to A,$$

is therefore an $F$-algebra isomorphism satisfying $h \circ \sigma = \sigma \circ h$. By [8, (3.1)], there exists a decomposition $(A,\sigma) \simeq \bigotimes_{i=1}^n (Q_i, \sigma_i)$ into quaternion $K$-algebras with orthogonal involution such that

$$\mathrm{disc}\, \sigma_i = \alpha_i K^{\times 2} \in K^\times / K^{\times 2}, \quad i = 1, \cdots, n.$$

Set $Q_i' = h(Q_i)$, $i = 1, \cdots, n$. We claim that every $Q_i'$ is a $\sigma$-invariant quaternion $K$-subalgebra of $A$. Let $(1, u_i, v_i, w_i)$ be a quaternion basis of $Q_i$ and set $u_i' = h(u_i)$, $v_i' = h(v_i)$ and $w_i' = h(w_i)$. Since $h(\alpha) = \iota(\alpha) \in K$ for every $\alpha \in K$, the condition $u_i^2 + u_i \in K$ implies that $u_i'^2 + u_i' \in K$. Similarly, as $h(\alpha) = \iota(\alpha) \in K^\times$ for every $\alpha \in K^\times$, we have $v_i'^2 \in K^\times$. The equality $u_i v_i + v_i u_i = v_i$ implies that $u_i' v_i' + v_i' u_i' = v_i'$. Hence, $Q_i'$ is a quaternion $K$-algebra with a quaternion basis $(1, u_i', v_i', w_i')$ for $i = 1, \cdots, n$. We also have

$$\sigma(Q_i') = \sigma(h(Q_i)) = h(\sigma(Q_i)) = h(Q_i) = Q_i',$$

proving the claim. Now, choose $x_i \in \mathrm{Alt}(Q_i, \sigma_i)$ with $x_i^2 = \alpha_i$, $i = 1, \cdots, n$. Then $h(x_i) \in \mathrm{Alt}(Q_i', \sigma|_{Q_i'})$ satisfies $h(x_i)^2 = \iota(\alpha_i)$, hence $\mathrm{disc}\, \sigma|_{Q_i'} = \iota(\alpha_i)K^{\times 2}$. We also have $(A, \sigma) \simeq \bigotimes_{i=1}^n (Q_i', \sigma|_{Q_i'})$, which implies in particular that $\mathfrak{Pf}(A, \sigma) \simeq \langle\langle \iota(\alpha_1), \cdots, \iota(\alpha_n) \rangle\rangle$. □

**Proposition 5.2.** *Let $(A, \sigma)$ be a totally decomposable algebra with orthogonal involution over $K$. If $(A, \sigma)$ has a descent to $F$, then there exists a totally singular conic $F$-algebra $S \subseteq \Phi(A, \sigma)$ for which $\Phi(A, \sigma) \simeq S \otimes_F K$ as $K$-algebras.*

*Proof.* Let $(A, \sigma) \simeq \bigotimes_{i=1}^n (Q_i, \sigma_i)$ be a decomposition of $(A, \sigma)$ into quaternion $K$-algebras with involution. Let $v_i \in \mathrm{Alt}(Q_i, \sigma_i)$ be a unit and set $\alpha_i = v_i^2 \in K^\times$, $i = 1, \cdots, n$. Then $\mathfrak{Pf}(A, \sigma) \simeq \langle\langle \alpha_1, \cdots, \alpha_n \rangle\rangle$ and $\Phi(A, \sigma) \simeq K[v_1, \cdots, v_n]$. Write $\alpha_i = b_i + c_i\eta$ for some $b_i, c_i \in F$, $i = 1, \cdots, n$. By (5.1) we have

$$\mathfrak{Pf}(A, \sigma) \simeq \langle\langle b_1 + c_1 + c_1\eta, \cdots, b_n + c_n + c_n\eta \rangle\rangle.$$

By [7, (5.6)], there exists $v_i' \in \mathrm{Alt}(A, \sigma)$ such that $v_i'^2 = b_i + c_i + c_i\eta$, $i = 1, \cdots, n$, and $\Phi(A, \sigma) \simeq K[v_1', \cdots, v_n']$. For $i = 1, \cdots, n$, we may identify $v_i$ (resp. $v_i'$) with an element of $\Phi(A, \sigma)$. Set

$$u_i = v_i + v_i' \in \Phi(A, \sigma) \quad \text{and} \quad w_i = (1 + \eta)v_i + \eta v_i' \in \Phi(A, \sigma),$$

so that $u_i^2 = c_i \in F$ and $w_i^2 = b_i + c_i\delta \in F$, $i = 1, \cdots, n$. Set $S = F[u_1, \cdots, u_n, w_1, \cdots, w_n]$. Then $S$ is a totally singular conic $F$-algebra and

$$S \otimes_F K \simeq K[u_1, \cdots, u_n, w_1, \cdots, w_n] \subseteq \Phi(A, \sigma).$$

On the other hand, as $v_i = \eta u_i + w_i$, the algebra $S \otimes_F K$ contains a copy of $v_i$ for $i = 1, \cdots, n$. Hence $\Phi(A, \sigma) \simeq S \otimes_F K$. □

**Remark 5.3.** According to [7, (3.2)], every totally singular conic $F$-algebra $S$ is a local algebra and its unique maximal ideal is $\mathfrak{m} = \{x \in S \mid x^2 = 0\}$. It follows that the unique maximal ideal of $S \otimes_F K$ is $\mathfrak{m} \otimes_F K$. In fact, every $x \in S \otimes_F K$ can be written as $x = u \otimes 1 + v \otimes \eta$ for some $u, v \in S$, so that $x^2 = (u^2 + \delta v^2) \otimes 1 + v^2 \otimes \eta$. Hence, $x^2 = 0$ if and only if $u^2 = v^2 = 0$, or equivalently, $x \in \mathfrak{m} \otimes_F K$.

**Lemma 5.4.** *Let $R$ be a totally singular conic algebra over $K$. Suppose that there exists a totally singular conic $F$-algebra $S$ such that $R \simeq S \otimes_F K$. If $E$ is a maximal subfield of $R$ containing $K$ and $L$ is a maximal subfield of $S$ containing $F$, then $E \simeq L \otimes_F K$.*

*Proof.* Let $\mathfrak{m}$ and $\mathfrak{m}'$ be the respective maximal ideals of $S$ and $R$. By (5.3), $\mathfrak{m} \otimes_F K$ is the unique maximal ideal of $S \otimes_F K$. Hence, the isomorphism $R \simeq S \otimes_F K$ induces a $K$-algebra isomorphism

$$R/\mathfrak{m}' \simeq (S \otimes_F K)/(\mathfrak{m} \otimes_F K).$$

Note that there exists a natural $K$-algebra isomorphism $(S \otimes_F K)/(\mathfrak{m} \otimes_F K) \simeq S/\mathfrak{m} \otimes_F K$, hence $R/\mathfrak{m}' \simeq S/\mathfrak{m} \otimes_F K$. By [7, (3.7 (i))], we have $E \simeq R/\mathfrak{m}'$ as $K$-algebras and $L \simeq S/\mathfrak{m}$ as $F$-algebras, hence $E \simeq L \otimes_F K$. □

**Lemma 5.5.** *Let $\mathfrak{b}$ be a bilinear Pfister form over a field $E$. If $\mathfrak{b}$ is isotropic, then there exist a positive integer $s$ and an anisotropic bilinear Pfister form $\mathfrak{c}$ over $E$ such that $\mathfrak{b} \simeq \langle\langle 1\rangle\rangle^s \otimes \mathfrak{c}$, where $\langle\langle 1\rangle\rangle^s$ is the $s$-fold Pfister form $\langle\langle 1, \cdots, 1\rangle\rangle$. Moreover, the integer $s$ is uniquely determined by the isomorphism class of $\mathfrak{b}$.*

*Proof.* See [1, p. 909]. □

**Notation 5.6.** We denote the integer $s$ in (5.5) by $\mathfrak{i}(\mathfrak{b})$. If $\mathfrak{b}$ is anisotropic, we set $\mathfrak{i}(\mathfrak{b}) = 0$. Also, if $(A, \sigma)$ is a totally decomposable algebra with orthogonal involution over $E$, we simply denote $\mathfrak{i}(\mathfrak{Pf}(A, \sigma))$ by $\mathfrak{i}(A, \sigma)$.

Note that [3, (7.5)] implies that $(A, \sigma)$ is anisotropic if and only if $\mathfrak{i}(A, \sigma) = 0$. Also, in view of [8, (3.1)], [3, (7.5)] and (4.6), if $\mathfrak{i}(A, \sigma) = s > 0$, then

$$(A, \sigma) \simeq (M_{2^s}(E), t) \otimes (B, \sigma'),$$

where $(B, \sigma')$ is a totally decomposable algebra with anisotropic orthogonal involution over $E$. Hence, $\Phi(A, \sigma) \simeq \Phi(M_{2^s}(E), t) \otimes \Phi(B, \sigma')$ by [7, (5.13)]. According to [7, (6.8)], $\Phi(B, \sigma')$ is a field. Also, [7, (5.7)] implies that $x^2 \in E^2$ for every $x \in \Phi(M_{2^s}(E), t)$. Hence, $\Phi(B, \sigma')$ may be identified with a maximal subfield of $\Phi(A, \sigma)$ containing $E$. It follows from [7, (3.7 (i))] that every maximal subfield of $\Phi(A, \sigma)$ containing $E$ is isomorphic to $\Phi(B, \sigma')$. In particular, such a maximal subfield has dimension $2^{n-s}$.

**Lemma 5.7.** *Let $(A, \sigma)$ be a totally decomposable algebra of degree $2^n$ with orthogonal involution over $K$ and let $r = n - \mathfrak{i}(A, \sigma)$. If $\Phi(A, \sigma) \simeq S \otimes_F K$ for some totally singular conic $F$-algebra $S$, then there exist $v_1, \cdots, v_n \in \Phi(A, \sigma)$ such that*

*(1) $\Phi(A, \sigma) = K[v_1, \cdots, v_n]$;*

*(2) $K[v_1, \cdots, v_r]$ is a maximal subfield of $\Phi(A, \sigma)$ with $v_i^2 \in F$ for $i = 1, \cdots, r$;*

*(3) $v_i^2 = 1$ for $i = r + 1, \cdots, n$.*

*Proof.* Let $L$ and $E$ be maximal subfields of $S$ and $\Phi(A, \sigma)$ respectively, with $F \subseteq L$ and $K \subseteq E$. By (5.4) we have $E \simeq L \otimes_F K$ as $K$-algebras. Also, as already observed, we have $\dim_F L = \dim_K E = 2^r$. Write $L = F[v_1, \cdots, v_r]$ for some $v_1, \cdots, v_r \in L$. Then $E \simeq K[v_1, \cdots, v_r]$. Since $\Phi(A, \sigma)$ is generated, as a $K$-algebra, by $n$ elements, by [7, (3.9)] there exist $v_{r+1}, \cdots, v_n \in \Phi(A, \sigma)$ such that $\Phi(A, \sigma) = K[v_1, \cdots, v_n]$ and $v_i^2 = 0$, $i = r+1, \cdots, n$. This proves parts (1) and (2). The third part follows by replacing $v_i$ with $v_i + 1$ for $i = r+1, \cdots, n$. □

# 6 Applications to the Pfister invariant

The following notation was used in [1].

**Notation 6.1.** For a bilinear form $\mathfrak{b}$ over a field $E$ we use the notation $Q(\mathfrak{b}) = \{\mathfrak{b}(v, v) \mid v \in V\}$, where $V$ is an underlying vector space of $\mathfrak{b}$.

Note that $Q(\mathfrak{b})$ is an $E^2$-subspace of $E$. Also, since every sum of squares in $E$ is a square, we have $Q(\langle\langle 1\rangle\rangle^s \otimes \mathfrak{b}) = Q(\mathfrak{b})$ for every positive integer $s$.

**Lemma 6.2.** *Let $\mathfrak{b}$ and $\mathfrak{b}'$ be two anisotropic bilinear $n$-fold Pfister forms over a field $E$ and let $s$ be a positive integer. Then $\langle\langle 1\rangle\rangle^s \otimes \mathfrak{b} \simeq \langle\langle 1\rangle\rangle^s \otimes \mathfrak{b}'$ if and only if $Q(\mathfrak{b}) = Q(\mathfrak{b}')$.*

*Proof.* See [1, p. 909]. □

We continue to assume that $K = F(\eta)$ is a separable quadratic extension of $F$ with $\eta^2 + \eta = \delta \in F \setminus \wp(F)$.

**Lemma 6.3.** *Let $(A, \sigma)$ be a totally decomposable algebra of degree $2^n$ with isotropic orthogonal involution over $K$. If $\Phi(A, \sigma) \simeq S \otimes_F K$ for some totally singular conic $F$-algebra $S$, then there exists a symmetric bilinear form $\mathfrak{b}$ over $F$ such that $\mathfrak{Pf}(A, \sigma) \simeq \mathfrak{b}_K$.*

*Proof.* Set $s = \mathfrak{i}(A, \sigma) > 0$ and $r = n - s$. By (5.7), one can write $\Phi(A, \sigma) = K[v_1, \cdots, v_n]$ for some $v_1, \cdots, v_n \in \Phi(A, \sigma)$ such that $K[v_1, \cdots, v_r]$ is a maximal subfield of $\Phi(A, \sigma)$ with $v_i^2 \in F^\times$, $i = 1, \cdots, r$, and $v_i^2 = 1$ for $i = r+1, \cdots, n$. Set $\alpha_i = v_i^2 \in F^\times$, $i = 1, \cdots, n$. Let $\mathfrak{b}$ be the bilinear Pfister form $\langle\langle \alpha_1, \cdots, \alpha_n \rangle\rangle$ over $F$. Then $\mathfrak{b} \simeq \langle\langle 1\rangle\rangle^s \otimes \mathfrak{c}$, where $\mathfrak{c} = \langle\langle \alpha_1, \cdots, \alpha_r \rangle\rangle$. The algebra $F[v_1, \cdots, v_n]$ may be considered as an underlying vector space of $\mathfrak{b}$ such that $\mathfrak{b}(x, x) = x^2$ for $x \in F[v_1, \cdots, v_n]$. Hence, $\Phi(A, \sigma) \simeq F[v_1, \cdots, v_n] \otimes_F K$ may be considered as an underlying $K$-vector space of $\mathfrak{b}_K$ such that

$$\mathfrak{b}_K(x, x) = x^2 \quad \text{for} \quad x \in \Phi(A, \sigma). \tag{1}$$

Since $K[v_1, \cdots, v_r]$ is a field, the form $\mathfrak{c}_K$ is anisotropic. On the other hand, there exists an anisotropic $r$-fold Pfister form $\mathfrak{c}'$ over $K$ for which $\mathfrak{Pf}(A, \sigma) \simeq \langle\langle 1\rangle\rangle^s \otimes \mathfrak{c}'$. Since $\mathfrak{Pf}(A, \sigma)(x, x) = x^2$ for every $x \in \Phi(A, \sigma)$, (1) implies that $Q(\mathfrak{b}_K) = Q(\mathfrak{Pf}(A, \sigma))$, i.e., $Q(\mathfrak{c}_K) = Q(\mathfrak{c}')$. Using (6.2) we get

$$\mathfrak{Pf}(A, \sigma) \simeq \langle\langle 1\rangle\rangle^s \otimes \mathfrak{c}' \simeq \langle\langle 1\rangle\rangle^s \otimes \mathfrak{c}_K \simeq \mathfrak{b}_K.$$ □

**Lemma 6.4.** *Let $(A, \sigma)$ be a central simple algebra with involution over a field $E$. For every $x \in \mathrm{Sym}(A, \sigma)$ and $y \in \mathrm{Alt}(A, \sigma)$ we have $xyx \in \mathrm{Alt}(A, \sigma)$.*

*Proof.* Write $y = z - \sigma(z)$ for some $z \in A$. Then

$$xyx = x(z - \sigma(z))x = xzx - \sigma(xzx) \in \mathrm{Alt}(A, \sigma).$$ □

The next result follows from [3, (6.1)].

**Lemma 6.5.** *Let $(A, \sigma)$ be a totally decomposable algebra with anisotropic orthogonal involution over a field $E$ and let $x \in A$. If $\sigma(x)x \in \mathrm{Alt}(A, \sigma)$, then $x = 0$.*

**Corollary 6.6.** *Let $(A, \sigma)$ be a totally decomposable algebra with anisotropic orthogonal involution over a field $E$ and let $x \in \mathrm{Sym}(A, \sigma)$. If $x^2 + \alpha \in \mathrm{Alt}(A, \sigma)$ for some $\alpha \in E$, then $x^2 = \alpha$.*

*Proof.* By (6.4) we have $x^4 + x^2\alpha = x(x^2 + \alpha)x \in \mathrm{Alt}(A, \sigma)$. Hence

$$\sigma(x^2 + \alpha) \cdot (x^2 + \alpha) = x^4 + \alpha^2 = x^4 + x^2\alpha + \alpha(x^2 + \alpha) \in \mathrm{Alt}(A, \sigma),$$

which implies that $x^2 = \alpha$ by (6.5). □

**Proposition 6.7.** *Let $(A,\sigma)$ be a totally decomposable algebra with orthogonal involution over $K$. If $(A,\sigma)$ has a descent to $F$, then there exists a symmetric bilinear form $\mathfrak{b}$ over $F$ such that $\mathfrak{Pf}(A,\sigma) \simeq \mathfrak{b}_K$.*

*Proof.* If $\sigma$ is isotropic, the result follows from (5.2) and (6.3). Suppose that $\sigma$ is anisotropic. Let $(B,\sigma')$ be a descent of $(A,\sigma)$ to $F$ and consider an isomorphism of $K$-algebras with involution

$$g : (A,\sigma) \xrightarrow{\sim} (B,\sigma') \otimes_F (K, \mathrm{id}).$$

Write $\Phi(A,\sigma) = K[v_1,\cdots,v_n]$ for some $v_1,\cdots,v_n \in \mathrm{Alt}(A,\sigma)$. As $g(\mathrm{Alt}(A,\sigma)) = \mathrm{Alt}(B,\sigma') \otimes_F K$, one can write $g(v_i) = u_i \otimes 1 + w_i \otimes \eta$ for some $u_i, w_i \in \mathrm{Alt}(B,\sigma')$, $i = 1,\cdots,n$. We first show that $u_i^2, w_i^2 \in F$ for every $i$. We have

$$g(v_i)^2 = (u_i \otimes 1 + w_i \otimes \eta)^2 = (u_i^2 + \delta w_i^2) \otimes 1 + (u_i w_i + w_i u_i + w_i^2) \otimes \eta. \quad (2)$$

Set $\alpha_i = u_i w_i + w_i u_i + w_i^2$. Then

$$w_i^2 + \alpha_i = u_i w_i + w_i u_i = u_i w_i - \sigma'(u_i w_i) \in \mathrm{Alt}(B,\sigma').$$

Since $g(v_i)^2 \in K$ we have $\alpha_i \in F$. As $\sigma$ is anisotropic, $\sigma'$ is also anisotropic. Hence (6.6) implies that

$$w_i^2 = \alpha_i \in F, \quad (3)$$

i.e., $u_i w_i = w_i u_i$. Since $g(v_i)^2 \in K$, (2) and (3) imply that $u_i^2 \in F$ for every $i$.

We now show that $w_i w_j = w_j w_i$ for $i,j = 1,\cdots,n$. As $(w_i + w_j)^2 = \alpha_i + \alpha_j + w_i w_j + w_j w_i$, we have $(w_i + w_j)^2 + \alpha_i + \alpha_j \in \mathrm{Alt}(B,\sigma')$. Again, (6.6) implies that $(w_i + w_j)^2 = \alpha_i + \alpha_j$, i.e., $w_i w_j = w_j w_i$. A similar argument shows that $w_i u_j = u_j w_i$ and $u_i u_j = u_j u_i$ for $i,j = 1,\cdots,n$.

Set $S = F[u_1,\cdots,u_n,w_1,\cdots,w_n]$. Then $S$ is a totally singular conic $F$-subalgebra of $B$. Since $\Phi(A,\sigma) = K[v_1,\cdots,v_n]$ and $v_i = g^{-1}(u_i \otimes 1 + w_i \otimes \eta)$ for every $i$, we have $\Phi(A,\sigma) \subseteq g^{-1}(S \otimes_F K)$. On the other hand $\Phi(A,\sigma)$ is maximal commutative and $g^{-1}(S \otimes_F K) \subseteq A$ is commutative. Hence $\Phi(A,\sigma) = g^{-1}(S \otimes_F K)$. We claim that

$$\mathfrak{Pf}(A,\sigma)(g^{-1}(s_1 \otimes 1), g^{-1}(s_2 \otimes 1)) \in F \quad \text{for} \quad s_1, s_2 \in S.$$

Write $\mathfrak{Pf}(A,\sigma)(g^{-1}(s_1 \otimes 1), g^{-1}(s_2 \otimes 1)) = a + b\eta$ for some $a,b \in F$. Then

$$g^{-1}(s_1 s_1 \otimes 1) + a + b\eta = g^{-1}(s_1 \otimes 1) g^{-1}(s_2 \otimes 1) + a + b\eta \in \mathrm{Alt}(A,\sigma).$$

The equality $g(\mathrm{Alt}(A,\sigma)) = \mathrm{Alt}(B,\sigma') \otimes_F K$ then implies that $(s_1 s_2 + a) \otimes 1 + b \otimes \eta \in \mathrm{Alt}(B,\sigma') \otimes_F K$, so $b \in \mathrm{Alt}(B,\sigma')$. Since $\sigma'$ is orthogonal, we have $1 \notin \mathrm{Alt}(B,\sigma')$, hence $b = 0$. This proves the claim. Finally, define the bilinear form $\mathfrak{b} : S \times S \to F$ via

$$\mathfrak{b}(s_1,s_2) = \mathfrak{Pf}(A,\sigma)(g^{-1}(s_1 \otimes 1), g^{-1}(s_2 \otimes 1)).$$

Then the restriction of $g^{-1}$ to $S \otimes_F K$ defines an isometry $(S \otimes_F K, \mathfrak{b}_K) \xrightarrow{\sim} (\Phi(A,\sigma), \mathfrak{Pf}(A,\sigma))$. □

# 7 Separable descent in general case

We continue to assume that $K = F(\eta)$, where $\eta^2 + \eta = \delta \in F \setminus \wp(F)$.

**Lemma 7.1.** *Let $E = K(\sqrt{\alpha})$ and $L = F(\sqrt{\alpha})$ for some $\alpha \in F \setminus K^2$. Let $A$ be a central simple algebra over $K$. If $\mathrm{cor}_{K/F}(A)$ splits, then $\mathrm{cor}_{E/L}(A \otimes_K E)$ is also split.*

*Proof.* If $\mathrm{cor}_{K/F}(A)$ splits, then by [5, (3.1 (2))] there exists an involution $\sigma$ of the second kind on $A$ with $\sigma|_F = \mathrm{id}$. Note that $E = L(\eta)$ is a separable quadratic extension of $L$. Let $\iota$ be the nontrivial automorphism of $E/L$. Then the map $\sigma \otimes \iota$ is an involution of the second kind on $A \otimes_K E$ which leaves $L$ elementwise invariant. Hence $\mathrm{cor}_{E/L}(A \otimes_K E)$ splits by [5, (3.1 (2))]. □

**Lemma 7.2.** *Let $(A, \sigma)$ be a central simple algebra with involution of the first kind over $K$. If $(A, \sigma)$ has a descent to $F$, then $\mathrm{cor}_{K/F}(A)$ splits.*

*Proof.* The result follows from [5, (3.13 (5))] and [5, (3.1 (1))]. □

**Theorem 7.3.** *For a totally decomposable algebra with orthogonal involution $(A, \sigma)$ over $K$ the following conditions are equivalent.*

*(1) $(A, \sigma)$ has a descent to $F$.*

*(2) $(A, \sigma)$ has a totally decomposable descent to $F$.*

*(3) $\mathrm{cor}_{K/F}(A)$ splits and $\mathfrak{Pf}(A, \sigma) \simeq \mathfrak{b}_K$ for some symmetric bilinear form $\mathfrak{b}$ over $F$.*

*Furthermore, if $\sigma$ is isotropic, the above conditions are equivalent to:*

*(4) $\mathrm{cor}_{K/F}(A)$ splits and $\Phi(A, \sigma) \simeq S \otimes_F K$ for some totally singular conic $F$-algebra $S$.*

*Proof.* The implication (2) ⇒ (1) is evident and (1) ⇒ (3) follows from (7.2) and (6.7).

(3) ⇒ (2): If $\mathfrak{Pf}(A, \sigma) \simeq \mathfrak{b}_K$ for some symmetric bilinear form $\mathfrak{b}$ over $F$, then by [3, (3.3)], $\mathfrak{b}$ is similar to a Pfister form $\mathfrak{c}$ over $F$. Hence $\mathfrak{c}_K$ is similar to $\mathfrak{Pf}(A, \sigma)$. Since $\mathfrak{c}_K$ and $\mathfrak{Pf}(A, \sigma)$ are both Pfister forms, we get $\mathfrak{c}_K \simeq \mathfrak{Pf}(A, \sigma)$ by [3, (2.4)]. Write $\mathfrak{c} = \langle\langle \alpha_1, \cdots, \alpha_n \rangle\rangle$ for some $\alpha_1, \cdots, \alpha_n \in F^\times$. Then $\mathfrak{Pf}(A, \sigma) \simeq \langle\langle \alpha_1, \cdots, \alpha_n \rangle\rangle_K$.

By [8, (3.1)] there exists a decomposition $(A, \sigma) \simeq \bigotimes_{i=1}^n (Q_i, \sigma_i)$ into quaternion algebras with orthogonal involution over $K$ such that $\mathrm{disc}\, \sigma_i = \alpha_i K^{\times 2} \in K^\times / K^{\times 2}$, $i = 1, \cdots, n$. If $\alpha_i \in F^{\times 2}$ for every $i$, then (4.6) shows that

$$(A, \sigma) \simeq_K \bigotimes_{i=1}^n (M_2(K), t) \simeq_K \bigotimes_{i=1}^n (M_2(F), t) \otimes_F (K, \mathrm{id}),$$

and we are done. Otherwise, (by re-indexing) we may assume that $\alpha_n \notin F^{\times 2}$. Choose $v_i \in \mathrm{Alt}(Q_i, \sigma_i)$ such that $v_i^2 = \alpha_i \in F^\times$, $i = 1, \cdots, n$. By induction on $n$ we prove a stronger result than we need, namely that for $i = 1, \cdots, n$, there exists a quaternion algebra with involution $(M_i, \tau_i)$ over $F$ such that

$$(A, \sigma) \simeq_K \bigotimes_{i=1}^n (M_i, \tau_i) \otimes_F (K, \mathrm{id}),$$

and $v_i \in \mathrm{Alt}(M_i, \tau_i)$. The case $n = 1$ follows from (4.7), hence let $n \geq 2$. Set $E = K[v_n]$ and $L = F[v_n] = F(\sqrt{\alpha_n})$. Since $K^{\times 2} \cap F^\times = F^{\times 2}$, we have $\alpha_n \notin K^{\times 2}$, so $E = K(\sqrt{\alpha_n})$. Let $B = C_A(v_n)$ be the centralizer of $v_n$ in $A$ (we have identified $v_n \in Q_n$ with an element of $A$). Then

$$(B, \sigma|_B) \simeq_E \bigotimes_{i=1}^{n-1} (Q_i, \sigma_i) \otimes_K (E, \mathrm{id}),$$

is a totally decomposable $E$-algebra of degree $2^{n-1}$ with orthogonal involution. By (7.1), $\mathrm{cor}_{E/L}(A \otimes_K E)$ splits. Since $B \sim A \otimes_K E$, by [9, Ch. 8, (9.8)] we have $\mathrm{cor}_{E/L}(B) \sim \mathrm{cor}_{E/L}(A \otimes_K E)$, hence $\mathrm{cor}_{E/L}(B)$ is also split. Note that $\mathfrak{Pf}(B, \sigma|_B) \simeq \langle\langle \alpha_1, \cdots, \alpha_{n-1} \rangle\rangle_E$ and $E = L(\eta)$ is a separable quadratic extension of $L$. By identifying $v_i \otimes 1 \in \mathrm{Alt}((Q_i, \sigma_i) \otimes_K (E, \mathrm{id}))$ with $v_i$, the induction hypothesis implies that for $i = 1, \cdots, n-1$, there exists a quaternion algebra with orthogonal involution $(Q_i', \sigma_i')$ over $L$ such that

$$(B, \sigma|_B) \simeq_E \bigotimes_{i=1}^{n-1} (Q_i', \sigma_i') \otimes_L (E, \mathrm{id}),$$

and $v_i \in \mathrm{Alt}(Q_i', \sigma_i')$. Since $L^2 \subseteq F$, by (3.3) there exists a quaternion $F$-algebra with orthogonal involution $(M_i, \tau_i)$ such that $(Q_i', \sigma_i') \simeq_L (M_i, \tau_i) \otimes_F (L, \mathrm{id})$ and $v_i \in \mathrm{Alt}(M_i, \tau_i)$, $i = 1, \cdots, n-1$. It follows that

$$(B, \sigma|_B) \simeq_E \bigotimes_{i=1}^{n-1} (M_i, \tau_i) \otimes_F (L, \mathrm{id}) \otimes_L (E, \mathrm{id}) \simeq_E \bigotimes_{i=1}^{n-1} (M_i, \tau_i) \otimes_F (E, \mathrm{id}).$$

The $K$-algebra $\bigotimes_{i=1}^{n-1} M_i \otimes_F K$ may be identified with a subalgebra of $A \subseteq B$. Set

$$Q_n'' = C_A(\textstyle\bigotimes_{i=1}^{n-1} M_i \otimes_F K) \quad \text{and} \quad \sigma_n'' = \sigma|_{Q_n''}.$$

Then $(Q_n'', \sigma_n'')$ is a quaternion $K$-algebra with orthogonal involution and

$$(A, \sigma) \simeq_K \bigotimes_{i=1}^{n-1} (M_i, \tau_i) \otimes_F (K, \mathrm{id}) \otimes_K (Q_n'', \sigma_n'').$$

Note that according to (4.1), $\mathrm{cor}_{K/F}(M_i \otimes_F K)$ splits for every $i$. Since $\mathrm{cor}_{K/F}(A)$ split, [5, (3.13 (2))] implies that $\mathrm{cor}_{K/F}(Q_n'')$ is also split. As $v_n$ commutes with $B$ and $\bigotimes_{i=1}^{n-1} M_i \otimes_F K \subseteq B$, we have $v_n \in Q_n''$. Using [6, (3.5)], we get $v_n \in \mathrm{Alt}(Q_n'', \sigma_n'')$. Hence (4.7) implies that $(Q_n'', \sigma_n'') \simeq_K (M_n, \tau_n) \otimes_F (K, \mathrm{id})$ for some quaternion $F$-algebra with involution $(M_n, \tau_n)$ with $v_n \in \mathrm{Alt}(M_n, \tau_n)$. It follows that

$$(A, \sigma) \simeq_K \bigotimes_{i=1}^{n} (M_i, \tau_i) \otimes_F (K, \mathrm{id}),$$

and $v_i \in \mathrm{Alt}(M_i, \tau_i)$, $i = 1, \cdots, n$. This completes the proof of (3) $\Rightarrow$ (2).

Finally, the implication (1) $\Rightarrow$ (4) follows from (7.2) and (5.2), even without the isotropy condition on $\sigma$. If $\sigma$ is isotropic, the implication (4) $\Rightarrow$ (3) follows from (6.3). $\square$

We conclude by an example which shows that if $\sigma$ is anisotropic, the implication (4) $\Rightarrow$ (1) in (7.3) does not hold.

**Example 7.4.** Suppose that $F^2 \neq F$. Let $\lambda \in F \setminus F^2$ and set $\beta = \lambda + \delta + \eta \in K$. Consider the involution $\sigma : M_2(K) \to M_2(K)$ defined by

$$\sigma \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} a & c\beta^{-1} \\ b\beta & d \end{pmatrix}.$$

Set

$$v = \begin{pmatrix} 0 & 1 \\ \beta & 0 \end{pmatrix} \quad \text{and} \quad w = \begin{pmatrix} \eta & 1 \\ \beta & \eta \end{pmatrix}.$$

Then $v \in \mathrm{Alt}(M_2(K), \sigma)$ and $w \in \mathrm{Sym}(M_2(K), \sigma)$. We also have $v^2 = \beta$ and $w = v + \eta$ (we have identified scalar matrices in $M_2(K)$ with elements of $K$). In particular, $\operatorname{disc}\sigma = \beta K^\times / K^{\times 2}$. Note that $\operatorname{cor}_{K/F}(M_2(K))$ splits and $\Phi(A, \sigma) = K[v] = K[w] \simeq F[w] \otimes_F K$. Since $w^2 = \lambda \in F$, $F[w]$ is a totally singular conic $F$-algebra. Hence, the pair $(M_2(K), \sigma)$ satisfies the condition (4) in (7.3). If $(M_2(K), \sigma)$ has a descent to $F$, then by (4.7), there exists $u \in \mathrm{Alt}(M_2(K), \sigma)$ such that $u^2 \in F^\times$. As $\mathrm{Alt}(M_2(K), \sigma)$ is one-dimensional, we have $\mathrm{Alt}(M_2(K), \sigma) = Kv$. Write $u = (b + c\eta)v$ for some $b, c \in F$. Then

$$u^2 = (\lambda b^2 + \delta b^2 + \lambda\delta c^2 + c^2\delta^2 + \delta c^2) + (b^2 + c^2 + \lambda c^2)\eta.$$

Since $u^2 \in F^\times$, we have $b^2 + c^2 + \lambda c^2 = 0$. If $c = 0$, then $b = 0$ which implies that $u = 0$, contradicting $u^2 \in F^\times$. If $c \neq 0$, then $\lambda = b^2 c^{-2} + 1 \in F^2$, which is again a contradiction. Hence, $(M_2(K), \sigma)$ has no descent to $F$, i.e., the implication (4) $\Rightarrow$ (1) in (7.3) does not hold for $(M_2(K), \sigma)$.

# References


[1] J. Arason, R. Baeza, Relations in $I^n$ and $I^n W_q$ in characteristic 2. *J. Algebra* **314** (2007), no. 2, 895–911.

[2] H. Dherte, Quadratic descent of involutions in degree 2 and 4. *Proc. Am. Math. Soc.* **123** (1995), no. 7, 1963–1969 .

[3] A. Dolphin, Orthogonal Pfister involutions in characteristic two. *J. Pure Appl. Algebra* **218** (2014), no. 10, 1900–1915.

[4] R. Elman, N. Karpenko, A. Merkurjev, *The algebraic and geometric theory of quadratic forms.* American Mathematical Society Colloquium Publications, 56. American Mathematical Society, Providence, RI, 2008.

[5] M.-A. Knus, A. S. Merkurjev, M. Rost, J.-P. Tignol, *The book of involutions.* American Mathematical Society Colloquium Publications, 44. American Mathematical Society, Providence, RI, 1998.

[6] M. G. Mahmoudi, A.-H. Nokhodkar, On split products of quaternion algebras with involution in characteristic two. *J. Pure Appl. Algebra* **218** (2014), no. 4, 731–734.

[7] M. G. Mahmoudi, A.-H. Nokhodkar, On totally decomposable algebras with involution in characteristic two. *J. Algebra* **451** (2016), 208–231.

[8] A.-H. Nokhodkar, The chain equivalence of totally decomposable orthogonal involutions in characteristic two. LAG preprint server, http://www.math.uni-bielefeld.de/LAG/man/569.html.

[9] W. Scharlau, *Quadratic and Hermitian forms.* Grundlehren der Mathematischen Wissenschaften, **270**. Springer-Verlag, Berlin, 1985.



A.-H. Nokhodkar, a.nokhodkar@kashanu.ac.ir,
Department of Pure Mathematics, Faculty of Science, University of Kashan, P. O. Box 87317-51167, Kashan, Iran.